

Nonparametric volatility density estimation for discrete time models

Bert van Es, Peter Spreij
Korteweg-de Vries Institute for Mathematics
Universiteit van Amsterdam
Plantage Muidergracht 24
1018 TV Amsterdam
The Netherlands

Harry van Zanten
Division of Mathematics and Computer Science
Vrije Universiteit
De Boelelaan 1081 a
1081 HV Amsterdam
The Netherlands

February 1, 2008

Abstract

We consider discrete time models for asset prices with a stationary volatility process. We aim at estimating the multivariate density of this process at a set of consecutive time instants.

A Fourier type deconvolution kernel density estimator based on the logarithm of the squared process is proposed to estimate the volatility density. Expansions of the bias and bounds on the variance are derived.

Key words: stochastic volatility models, density estimation, kernel estimator, deconvolution, mixing

AMS subject classification: 62G07, 62M07, 62P20

1 Introduction

Suppose that we have price data $S_0, S_1, \dots$ of a certain asset in a financial market. Let X be the log-return process, defined by $X_t = \log S_t - \log S_{t-1}$. It is commonly believed that stochastic volatility models of the form

$$X_t = \sigma_t Z_t \tag{1.1}$$

describe much of the observed behaviour of this type of data. Here Z is typically an i.i.d. noise sequence (often Gaussian) and at each time t the random variables σ_t and Z_t are independent. We will assume that the process σ is strictly stationary and that the (multivariate) marginal distributions of σ have a density with respect to the Lebesgue measure on $(0, \infty)$. Our aim is to construct a nonparametric estimator for the multivariate density of $(\sigma_t, \dots, \sigma_{t+p})$, and to study its asymptotic behaviour.

Models that are used in the literature to describe the volatility display rather different invariant distributions. This observation lies at the basis of our point of view, which we pursue in this paper, that nonparametric estimation procedures are by all means sensible tools to get some insight in the behaviour of the volatility. Quite often in models that are used in practice, the invariant distributions of σ are unimodal. Since it is known that volatility clustering is an often occurring phenomenon, it is hard to believe that this can be explained by any of these models. Instead, one would expect in such a case for instance the distribution of (σ_t, σ_{t+1}) to have a density that has concentration regions around the diagonal with possibly peaks at certain clusters of low and high volatility, a phenomenon that may lead to for instance bimodal one-dimensional marginal distributions. Nonparametric density estimation could perhaps reveal such a shape of the invariant density of the volatility.

We will distinguish two classes of models in this paper. In both of them we will assume that the noise sequence is standard Gaussian and that σ is a strictly stationary, positive process satisfying a certain mixing condition. The way in which the bivariate process (σ, Z) , in particular its dependence structure, is further modelled differs however. In the first class of models that we consider, we assume that the process σ is predictable with respect to the filtration $\mathcal{F}_t$ generated by the process Z . Note that σ_t is independent of Z_t for each fixed time t . We furthermore have that (assuming that the unconditional variances are finite) σ_t^2 is equal to the conditional variance of X_t given $\mathcal{F}_{t-1}$. This class of models has become quite popular in the econometrics literature. Financial data such as log-returns of stock prices or exchange rates are believed to share a number of stylized features, including for instance heavy-tailedness and long-range dependence. Models of the type (1.1) have been proposed to capture those features. A well-known family included in the class (1.1) is the family of GARCH-models, introduced by Bollerslev (1986). For the GARCH(p, q)-model the sequence $\{\sigma_t\}$ in (1.1) is

assumed to satisfy the equation

$$\sigma_t^2 = \alpha_0 + \sum_{i=1}^p \alpha_i X_{t-i}^2 + \sum_{j=1}^q \beta_j \sigma_{t-j}^2, \quad (1.2)$$

where the α_i and β_j are nonnegative constants. Under suitable assumptions, see Bougerol and Picard (1992), GARCH processes are stationary and the statistical problem in this case would be to estimate the coefficients α_i and β_j in (1.2).

In the second class of models that we consider, we assume that the whole process σ is independent of the noise process Z . In this case, the natural underlying filtration $\mathcal{F} = \{\mathcal{F}_t\}_{t \geq 0}$ is generated by the two processes Z and σ in the following way. For each t the σ -algebra $\mathcal{F}_t$ is generated by Z_s , $s \leq t$ and σ_s , $s \leq t+1$. This choice of the filtration enforces σ to be predictable. As in the first model the process X becomes a martingale difference sequence and we have again (assuming that the unconditional variances are finite) that σ_t^2 is the conditional variance of X_t given $\mathcal{F}_{t-1}$. An example of such a model is given in De Vries (1991), where σ is generated as an AR(1) process with α -stable noise ($\alpha \in (0, 1)$).

As we said before, we do not want to make a parametric assumption such as (1.2), but we still want to measure the volatility of the data somehow. In the present paper we propose a nonparametric statistical procedure for this problem. Using ideas from deconvolution theory, we will propose a procedure for the estimation of the marginal density at a fixed point. To assess the quality of our procedure, we will derive expansions of the bias and bounds on the variance. This will be done separately for the two kinds of model classes outlined above.

2 Primer on kernel type deconvolution

We briefly review the construction of the deconvolution kernel density estimator based on i.i.d. observations, see also Wand and Jones (1995). For simplicity we consider in this section the univariate case only. Recall that the *characteristic function* or *Fourier transform* of a density function g is defined by

$$\phi_g(t) = \mathbb{E} e^{itX} = \int_{-\infty}^{\infty} e^{itx} g(x) dx, \quad (2.1)$$

where X is a random variable with density function g . In the standard deconvolution setting the random variable X is equal to the sum of two independent random variables, say Y , with unknown density f , and Z , with known density k . So g is the convolution of f and k and

$$\phi_g(t) = \mathbb{E} e^{itX} = \mathbb{E} e^{it(Y+Z)} = \mathbb{E} e^{itY} \mathbb{E} e^{itZ} = \phi_f(t) \phi_k(t). \quad (2.2)$$

The objective is to estimate f from i.i.d. observations of $X_1, \dots, X_n$ having density g . In identity (2.2) we know $\phi_k(t)$ and we can estimate $\phi_g(t)$ by the characteristic function of a *kernel estimator* g_{nh} of g . So

$$g_{nh}(x) = \frac{1}{n} \sum_{j=1}^n \frac{1}{h} w\left(\frac{x - X_j}{h}\right), \quad (2.3)$$

where w is an integrable function with integral one, called the *kernel function*, and $h > 0$ is a positive number, called the *bandwidth*, governing the curvature of the estimate. The kernel estimator itself is also a convolution of the empirical distribution function G_n of the observations and the rescaled kernel function $w_h(x) = w(x/h)/h$. So, with ϕ_w the Fourier transform of w ,

$$\phi_{g_{nh}}(t) = \phi_{w_h}(t) \int_{-\infty}^{\infty} e^{itx} dG_n(x) = \phi_w(ht) \phi_{emp}(t), \quad (2.4)$$

where

$$\phi_{emp}(t) = \int_{-\infty}^{\infty} e^{itx} dG_n(x) = \frac{1}{n} \sum_{j=1}^n e^{itX_j} \quad (2.5)$$

is called the *empirical characteristic function*. From (2.2) we see that

$$\frac{\phi_w(ht) \phi_{emp}(t)}{\phi_k(t)} \quad (2.6)$$

is an obvious candidate to estimate ϕ_f . Applying an inverse Fourier transform we obtain an estimator of f . Define the estimator f_{nh} of f as

$$f_{nh}(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \frac{\phi_w(ht) \phi_{emp}(t)}{\phi_k(t)} dt. \quad (2.7)$$

The inversion is allowed if the function (2.6) is integrable. In general this is not guaranteed. However, to enforce integrability, we assume that ϕ_w has a bounded support. Note that (2.7) can be rewritten as

$$\begin{aligned} f_{nh}(x) &= \frac{1}{n} \sum_{j=1}^n \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\phi_w(ht)}{\phi_k(t)} e^{-it(x-X_j)} dt \\ &= \frac{1}{nh} \sum_{j=1}^n \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\phi_w(s)}{\phi_k(s/h)} e^{-is(x-X_j)/h} ds \\ &= \frac{1}{nh} \sum_{j=1}^n v_h\left(\frac{x - X_j}{h}\right), \end{aligned} \quad (2.8)$$

where

$$v_h(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\phi_w(s)}{\phi_k(s/h)} e^{-isx} ds. \quad (2.9)$$

It is easy to see that the function v_h , and hence the estimator $f_{nh}(x)$, is real valued. Indeed, taking complex conjugates, we get

$$\begin{aligned}\overline{v_h(x)} &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{isx} \overline{\phi_w(s)/\phi_k(s/h)} ds \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} \overline{\phi_w(-s)/\phi_k(-s/h)} ds \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} \phi_w(s)/\phi_k(s/h) ds \\ &= v_h(x).\end{aligned}$$

A popular performance measure for deconvolution kernel estimators is the *mean squared error* (MSE). The MSE of $f_{nh}(x)$ is defined as $E(f_{nh}(x) - f(x))^2$. To obtain asymptotic expansions for the MSE, we need expansions for the bias and variance of the estimator. The expectation of $f_{nh}(x)$ is equal to the expectation of an ordinary kernel density estimator of f based on observations from f . We have

$$\begin{aligned}E f_{nh}(x) &= \int \frac{1}{h} w\left(\frac{x-u}{h}\right) f(u) du \\ &= f(x) + \frac{1}{2}h^2 \int u^2 w(u) du f''(x) + o(h^2),\end{aligned}$$

as $n \rightarrow \infty$, $h \rightarrow 0$ and $nh \rightarrow \infty$, provided that w is symmetric and f satisfies some smoothness conditions, essentially twice differentiability at x . The asymptotic variance of $f_{nh}(x)$ depends on the tails of the characteristic function of the density k . The smoother k , the faster the tails of the characteristic function vanish and the larger the asymptotic variance, see for instance Fan (1991).

3 Construction of the estimators

We consider the model (1.1), so $X_t = \sigma_t Z_t$. If we square this equation and take logarithms we get

$$\log X_t^2 = \log \sigma_t^2 + \log Z_t^2. \quad (3.1)$$

Recall that under our assumptions for each t the random variables σ_t and Z_t are independent. The density of $\log Z_t^2$, denoted by k , is given by

$$k(x) = \frac{1}{\sqrt{2\pi}} e^{\frac{1}{2}x} e^{-\frac{1}{2}e^x}. \quad (3.2)$$

Its graph is given in Figure 1 below.

As in Section 2, it seems reasonable to use a *deconvolution kernel density estimator* to estimate the unknown density f of $\log \sigma_t^2$. An estimate of




Figure 1: The density function k of $\log Z_t^2$.

the density of σ_t^2 or σ_t can then be obtained by a simple transformation. Computing the characteristic function ϕ_k of $\log Z^2$ we get, with $k(x)$ as in (3.2),

$$\phi_k(t) = \int_{-\infty}^{\infty} e^{itx} k(x) dx = \frac{1}{\sqrt{\pi}} 2^{it} \Gamma(\frac{1}{2} + it), \quad (3.3)$$

where the gamma function Γ is defined for all complex z with positive real part by

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt.$$

The graphs of $\text{Re}(\phi_k)$, $\text{Im}(\phi_k)$ and $|\phi_k|$ are given in Figures 2 and 3.




Figure 2: The real and imaginary part of the characteristic function ϕ_k .

For the model (1.1) this leads to the estimator

$$f_{nh}(x) = \frac{1}{nh} \sum_{j=1}^n v_h\left(\frac{x - \log X_j^2}{h}\right) \quad (3.4)$$




Figure 3: The modulus of ϕ_k .

of the density f of $\log \sigma_t^2$, with $v_h(x)$ as in (2.9). Note that, like in the previous section, this estimator is real valued.

The expression for the estimator of the density of the p -dimensional random vector $(\log \sigma_t^2, \dots, \log \sigma_{t-p+1}^2)$ is similar. We first introduce some auxiliary notation. Let p be fixed and write $\mathbf{x}_j$ for a vector $(x_j, \dots, x_{j-p+1})$. We use similar boldface expressions for other (random) vectors. The kernel $\mathbf{w}$ that we will use in the multivariate case is just a product kernel, $\mathbf{w}(\mathbf{x}) = \prod_{j=1}^p w(x_j)$. Likewise $\mathbf{k}(\mathbf{x}) = \prod_{j=1}^p k(x_j)$. Then with $\mathbf{v}_h$ defined by

$$\mathbf{v}_h(\mathbf{x}) = \frac{1}{(2\pi)^p} \int_{\mathbb{R}^p} \frac{\phi_{\mathbf{w}}(\mathbf{s})}{\phi_{\mathbf{k}}(\mathbf{s}/h)} e^{-i\mathbf{s} \cdot \mathbf{x}} d\mathbf{s}, \quad (3.5)$$

where $\mathbf{s} \in \mathbb{R}^p$ and $\cdot$ denotes inner product, the multivariate density estimator is given by

$$f_{nh}(\mathbf{x}) = \frac{1}{(n-p+1)h^p} \sum_{j=p}^n \mathbf{v}_h\left(\frac{\mathbf{x} - \log \mathbf{X}_j^2}{h}\right), \quad (3.6)$$

where we use $\log \mathbf{X}_j^2$ to denote the vector $(\log X_j^2, \dots, \log X_{j-p+1}^2)$.

4 Asymptotics

The bias of the deconvolution estimator described in Section 2 will be seen to be the same as the bias of a kernel density estimator based on independent observations from f . Hence, under standard smoothness assumptions, it is of order h^2 as $h \rightarrow 0$. The variance of this type of deconvolution estimator heavily depends on the rate of decay to zero of $|\phi_k(t)|$ as $|t| \rightarrow \infty$. The faster the decay the larger the asymptotic variance. In other words, the smoother k the harder the estimation problem. This follows for instance for i.i.d. observations from results in Fan (1991) and for stationary observations from the work of Masry (1991, 1993a,b).

The rate of decay of $|\phi_k(t)|$ for the density (3.2) is given by Lemma 5.1 in Section 5, where we show that

$$|\phi_k(t)| \sim \sqrt{2} e^{-\frac{1}{2}\pi|t|}, \quad \text{as } |t| \rightarrow \infty. \quad (4.1)$$

By the similarity of the tail of this characteristic function to the tail of a Cauchy characteristic function we can expect the same order of the mean squared error as in Cauchy deconvolution problems, where it decreases logarithmically in n , cf. Fan (1991) for results on i.i.d. observations. Note that this rate, however slow, is faster than the one for normal deconvolution.

In the model (3.1) the sequence $\{\log X_t^2\}$ is not independent, so results on the asymptotic behavior of the kernel estimator of Section 2 are not directly applicable. In the literature also more general deconvolution problems have been studied, where the i.i.d. assumption has been relaxed. For instance, the deconvolution model $X_j = Y_j + Z_j$, where $\{Y_j, Z_j\}$ is a stationary sequence and the sequences $\{Z_j\}$ $\{Y_j\}$ are independent has been treated by E. Masry (1991, 1993a,b).

Expansions for the variance of the deconvolution kernel estimator have been derived under several mixing conditions. Under the assumption that the volatility process is independent of the noise sequence, the model (3.1) fits into this scheme. We will obtain similar results for the estimator when σ (as a process) is not independent of Z , but only predictable with respect to the filtration generated by Z .

Let us define the mixing conditions. For a certain process $\{X_j\}$ let $\mathcal{F}_a^b$ be the σ -algebra of events generated by the random variables X_j , $j = a, \dots, b$. Let the mixing coefficient α_k be defined by

$$\alpha_k = \sup_{A \in \mathcal{F}_{-\infty}^0, B \in \mathcal{F}_k^\infty} |P(AB) - P(A)P(B)|. \quad (4.2)$$

We call a process $\{X_j\}$ *strongly mixing* if $\alpha_k \rightarrow 0$ as $k \rightarrow \infty$.

To obtain expansions for the bias and variance we also need conditions on the kernel function w such as bounded support of its characteristic function $\phi_w(t)$. Moreover, the rate of decay to zero of $\phi_w(t)$ at the boundary of its support turns up in the asymptotics. The complete list of assumptions on w that we use is the following.

Condition W. Let w be a real symmetric function satisfying

1. $\int_{-\infty}^{\infty} |w(u)| du < \infty$,
2. $\int_{-\infty}^{\infty} w(u) du = 1$,
3. $\int_{-\infty}^{\infty} u^2 w(u) du < \infty$,
4. $\lim_{|u| \rightarrow \infty} w(u) = 0$,

5. ϕ_w , the characteristic function of w has support $[-1,1]$,
6. $\phi_w(1-t) = At^\alpha + o(t^\alpha)$, as $t \downarrow 0$ for some $\alpha > 0$.

Note that by Fourier inversion these conditions imply that w is bounded and Lipschitz. More precisely, we have

$$|w(x)| \leq \frac{1}{2\pi} \quad \text{and} \quad |w(x+u) - w(x)| \leq \frac{1}{2\pi}|u|. \quad (4.3)$$

An example of such a kernel, from Wand (1998), with $\alpha = 3$ and $A = 8$, is

$$w(x) = \frac{48x(x^2 - 15) \cos x - 144(2x^2 - 5) \sin x}{\pi x^7}. \quad (4.4)$$

It has characteristic function

$$\phi_w(t) = (1 - t^2)^3, \quad |t| \leq 1. \quad (4.5)$$

The next theorem, whose proof can be found in Section 5, establishes the expansion of the bias and an order bound on the variance of our estimator under a strong mixing condition. Under broad conditions this mixing condition is satisfied if the process σ is a Markov chain, since then convergence of α_k to zero takes place at an *exponential rate*, see Theorems 4.2 and Theorem 4.3 of Bradley (1985) for precise statements. Similar behaviour occurs for ARMA processes with absolutely continuous distributions of the noise terms (Bradley (1985), Example 6.1).

Theorem 4.1. *Assume that the process X is strongly mixing with coefficient α_k satisfying*

$$\sum_{j=1}^{\infty} \alpha_j^\beta < \infty,$$

for some $\beta \in (0,1)$. Let the kernel function w satisfy Condition W and let the density f of the p -vector $(\log \sigma_1^2, \dots, \log \sigma_p^2)$ be bounded and twice continuously differentiable with bounded second order partial derivatives. Assume that σ is a predictable process with respect to the filtration generated by the process Z . Then we have for the estimator of the multivariate density defined as in (3.6) and $h \rightarrow 0$

$$\mathbb{E} f_{nh}(\mathbf{x}) = f(\mathbf{x}) + \frac{1}{2}h^2 \int \mathbf{u}^\top \nabla^2 f(\mathbf{x}) \mathbf{u} w(\mathbf{u}) d\mathbf{u} + o(h^2) \quad (4.6)$$

and

$$\text{Var} f_{nh}(\mathbf{x}) = O\left(\frac{1}{n} (h^{2\alpha-\beta} e^{\pi/h})^p\right). \quad (4.7)$$

Theorem 4.2. Assume that the process σ is strongly mixing with coefficient α_k satisfying

$$\sum_{j=1}^{\infty} \alpha_j^\beta < \infty,$$

for some $\beta \in (0, 1)$. Let the kernel function w satisfy Condition W and let the density f of the p -vector $(\log \sigma_1^2, \dots, \log \sigma_p^2)$ be bounded and twice continuously differentiable with bounded second order partial derivatives. Assume furthermore that σ and Z are independent processes. Then the multivariate density estimator f_{nh} satisfies the same bias expansion as in Theorem 4.1. For the variance we have the sharper bound

$$\text{Var } f_{nh}(\mathbf{x}) = O\left(\frac{1}{n} (h^{2\alpha} e^{\pi/h})^p\right). \quad (4.8)$$

Remark 4.3. Because of the exponential factor in the variance bound, in order to obtain consistency, one has to take essentially $h \geq \pi/\log n$, see also Stefanski (1990) for a related problem. On the other hand we would like to minimize the bias, so the choice $h = \pi/\log n$ is optimal. Both bias and variance decay at a logarithmic rate for this choice of bandwidth. This seems disappointing, however Fan (1991) shows for the i.i.d. situation of Section 2 that we can not expect anything better.

Remark 4.4. Notice that the results in Masry (1993a,b) establishing strong consistency, rates of convergence and asymptotic normality are not useful here, because the condition that ϕ_k has either purely real or purely imaginary tails is not satisfied.

Remark 4.5. Note that our assumptions in Theorem 4.1 are slightly different from those of Masry (1991). One of the essential facts that are used in the proof is the mixing property of X . If σ and Z are independent processes this is implied by a similar assumption on the σ process itself as in Masry (1991).

Remark 4.6. In the case where the processes σ and Z are independent, the estimators $f_{nh}(x)$ have the following property.

$$\tilde{f}_{nh}(x) := \mathbb{E}[f_{nh}(x) | \mathcal{F}^\sigma] = \frac{1}{nh} \sum_{j=1}^n w\left(\frac{x - \log \sigma_j^2}{h}\right), \quad (4.9)$$

where $\mathcal{F}^\sigma$ denotes the σ -algebra generated by the whole process σ . Thus the $\tilde{f}_{nh}(x)$ would be ordinary kernel density estimators, if the σ_j^2 could be observed.

Equation (4.9) is seen to be true as follows. Write $u_j = \log X_j^2$ and use similar notation for $\zeta_j = \log Z_j^2$ and $\tau_j = \log \sigma_j^2$. Then

$$\begin{aligned} \mathbb{E}[v_h(\frac{x - u_j}{h})|\mathcal{F}^\sigma] &= \frac{1}{2\pi} \int \mathbb{E} e^{is\zeta_j/h} \frac{\phi_w(s)}{\phi_k(s/h)} e^{-is(x-\tau_j)/h} ds \\ &= \frac{1}{2\pi} \int \phi_w(s) e^{-is(x-\tau_j)/h} ds \\ &= w(\frac{x - \tau_j}{h}). \end{aligned}$$

The result now follows. Of course, the analogous statement for the multivariate density estimator is equally true. One has

$$\tilde{f}_{nh}(\mathbf{x}) := \mathbb{E}[f_{nh}(\mathbf{x})|\mathcal{F}^\sigma] = \frac{1}{nh^p} \sum_{j=p}^n w\left(\frac{\mathbf{x} - (\log \sigma_j^2, \dots, \log \sigma_{j-p+1}^2)}{h}\right), \quad (4.10)$$

Remark 4.7. Better bounds on the asymptotic variance than in Theorem 4.1 can be obtained under stronger mixing conditions. Consider for instance *uniform mixing*. In this case the mixing coefficient ϕ_t is defined for $t > 0$ as

$$\phi_t = \sup_{A \in \mathcal{F}_{-\infty}^0, B \in \mathcal{F}_t^\infty} |P(A|B) - P(A)|. \quad (4.11)$$

Similar to strong mixing, a process is called *uniform mixing* if $\phi_t \rightarrow 0$ for $t \rightarrow \infty$. Obviously, uniform mixing implies strong mixing. As a matter of fact, one has the relation

$$\alpha_t \leq \frac{1}{2}\phi_t.$$

See Doukhan (1994) for this inequality and many other mixing properties. If $\{\sigma_t\}$ is uniform mixing with coefficient ϕ satisfying $\sum_{j=1}^\infty \phi(j)^{1/2} < \infty$, then the variance bound (4.7) can be replaced with

$$\text{Var } f_{nh}(\mathbf{x}) = O\left(\frac{1}{n} (h^{2\alpha} e^{\pi/h})^p\right). \quad (4.12)$$

The proof of the latter bound runs similarly to the strong-mixing bound as given in section 5. The essential difference is that in equation (5.5) we use Theorem 17.2.3 of Ibragimov and Linnik (1971) with $\tau = 0$ instead of Deo's (1973) lemma, as in the proof of Theorem 2 in Masry (1983). The result is that we can now bound the term M_{nh} of equation (5.5) by a constant times $\sum_{j=1}^{n-p+1} \varphi_j^{1/2} \mathbb{E} W_0^2$. After this step the proof is essentially unchanged. Use the estimate $\mathbb{E} W_0^2 \leq Ch^p \|v\|_2^2$ to finish the proof. Notice that this bound on the variance is of the same order as the one we obtained in Theorem 4.2, where σ was only assumed to be strongly mixing. This bound cannot be improved upon by strengthening the assumption to uniform mixing.

Remark 4.8. An example of an observed process that is stongly mixing and that belong to the first model class is a GARCH(p, q) process. It has been shown in Carasso and Chen (2002) (see also Boussama (1998)) that such a process is β -mixing with exponentially decaying β -mixing coefficients. Hence this process is also α -mixing, since the β -mixing coefficient $\beta_k = \text{E ess sup}\{|P(A|\mathcal{F}_k^\infty) - P(A)| : A \in \mathcal{F}_{-\infty}^0\}$ satisfies the inequality $2\alpha_k \leq \beta_k$ (see Doukhan (1994)). Notice that we also have that the assumption of Theorem 4.1 on the α 's is satisfied in this case.

5 Proofs

All the estimators that we proposed involve the functions ϕ_k and ϕ_w . For these functions and related ones we need expansions and order estimates. These are collected in the lemmas of this subsection.

Lemma 5.1. *For $|t| \rightarrow \infty$ we have*

$$\begin{aligned} |\phi_k(t)| &= \sqrt{2} e^{-\frac{1}{2}\pi|t|} (1 + O(\frac{1}{|t|})), \\ \text{Re } \phi_k(t) &= |\phi_k(t)| [\cos(t \log(\sqrt{1+4t^2} - t)) + O(\frac{1}{|t|})], \\ \text{Im } \phi_k(t) &= |\phi_k(t)| [\sin(t \log(\sqrt{1+4t^2} - t)) + O(\frac{1}{|t|})]. \end{aligned}$$

Proof. By the Stirling formula for the complex gamma function, cf. Abramowitz and Stegun (1964) Chapter 6, we have

$$\log \Gamma(z) = (z - \frac{1}{2}) \log z - z + \frac{1}{2} \log 2\pi + O(\frac{1}{|z|}), \quad (5.1)$$

as $|z| \rightarrow \infty$ and $|\text{Arg } z| < \pi$ for some $\delta > 0$. So for $z = \frac{1}{2} + it$ and $|t| \rightarrow \infty$ we get

$$\begin{aligned} \log \Gamma(\frac{1}{2} + it) &= it \log(\frac{1}{2} + it) - (\frac{1}{2} + it) + \frac{1}{2} \log 2\pi + O(\frac{1}{|t|}) \\ &= it(\log |\frac{1}{2} + it| + i \text{Arg}(\frac{1}{2} + it)) - (\frac{1}{2} + it) + \frac{1}{2} \log 2\pi + O(\frac{1}{|t|}) \\ &= -t \text{Arg}(\frac{1}{2} + it) - \frac{1}{2} + \frac{1}{2} \log 2\pi + i(t \log |\frac{1}{2} + it| - t) + O(\frac{1}{|t|}). \end{aligned}$$

Taking the modulus of the exponent the imaginary part vanishes and we get

$$\begin{aligned} |\Gamma(\frac{1}{2} + it)| &= \exp(-t \text{Arg}(\frac{1}{2} + it) - \frac{1}{2} + \frac{1}{2} \log 2\pi + O(\frac{1}{|t|})) \\ &= \sqrt{2\pi} \exp(-t \arctan 2t - \frac{1}{2} + O(\frac{1}{|t|})) \\ &= \sqrt{2\pi} \exp(-\frac{1}{2}\pi|t| + O(\frac{1}{|t|})) \\ &= \sqrt{2\pi} \exp(-\frac{1}{2}\pi|t|) (1 + O(\frac{1}{|t|})). \end{aligned}$$

Here we have used the expansion $t \arctan t = t(\frac{1}{2}\pi - \arctan(1/t)) = \frac{1}{2}\pi t - 1 + O(1/t)$, as t tends to infinity. For negative t a similar expansion holds. Since

$2^{it} = \exp(it \log 2)$ has modulus one, substituting this expansion in (3.3) now proves the first statement of the lemma. The argument of $\Gamma(\frac{1}{2} + it)$ satisfies

$$\begin{aligned} \text{Arg}(\Gamma(\tfrac{1}{2} + it)) &= t \log |\tfrac{1}{2} + it| - t + O(\tfrac{1}{|t|}) \\ &= -t \log 2 + t \log |1 + 2it| - t + O(\tfrac{1}{|t|}). \end{aligned}$$

So, since $\text{Arg}(2^{it}) = t \log 2$, we have

$$\text{Arg}(\phi_k(t)) = t \log(\sqrt{1 + 4t^2} - t) + O(\tfrac{1}{|t|}),$$

which proves the second and third statement of the lemma. $\square$

Consider now the function v_h defined in (2.9).

Lemma 5.2. *We have the following order estimate for the L^2 norm of v_h . For $h \rightarrow 0$*

$$\|v_h\|_2 = O(h^{\frac{1}{2} + \alpha} e^{\pi/2h}). \quad (5.2)$$

Proof. By Parseval's identity

$$\|v_h\|^2 = \frac{1}{2\pi} \int_{-1}^1 \left| \frac{\phi_w(s)}{\phi_k(s/h)} \right|^2 dx.$$

Write

$$\begin{aligned} & \int_{-1}^1 \left| \frac{\phi_w(s)}{\phi_k(s/h)} \right|^2 ds \\ & \leq \frac{1}{2} \int_{-1}^1 |\phi_w(s)|^2 e^{\pi|s/h|} ds \end{aligned} \quad (5.3)$$

$$+ \int_{-1}^1 |\phi_w(s)|^2 \left| \frac{1}{|\phi_k(s/h)|^2} - \frac{1}{2} e^{\pi|s/h|} \right| ds \quad (5.4)$$

The integral in (5.3) can be rewritten as

$$\begin{aligned} & \int_{-1}^1 |\phi_w(s)|^2 e^{\pi|s/h|} ds \\ & = e^{\pi/h} \int_{-1}^1 |\phi_w(s)|^2 e^{\pi(|s/h| - (1/h))} ds \\ & = 2e^{\pi/h} \int_0^1 |\phi_w(s)|^2 e^{\pi(|s/h| - (1/h))} ds \\ & = 2e^{\pi/h} h \int_0^{1/h} |\phi_w(1 - hv)|^2 e^{\pi((1 - hv)/h - (1/h))} dv \\ & = 2e^{\pi/h} h^{1+2\alpha} \int_0^{1/h} \left| \frac{\phi_w(1 - hv)}{(hv)^\alpha} \right|^2 v^{2\alpha} e^{-\pi v} dv \\ & \sim 2e^{\pi/h} h^{1+2\alpha} A^2 \int_0^\infty v^{2\alpha} e^{-\pi v} dv \\ & = 2e^{\pi/h} h^{1+2\alpha} (\pi)^{-1-2\alpha} A^2 \Gamma(2\alpha + 1), \end{aligned}$$

by the dominated convergence theorem. Omitting constants, we can rewrite the integral (5.4) as

$$\begin{aligned}
& \int_{-1}^1 |\phi_w(s)|^2 e^{\pi|s/h|} \left| \frac{2e^{-\pi|s/h|}}{|\phi_k(s/h)|^2} - 1 \right| ds \\
&= e^{\pi/h} \int_{-1}^1 |\phi_w(s)|^2 \left| \frac{2e^{-\pi|s/h|}}{|\phi_k(s/h)|^2} - 1 \right| e^{\pi(|s/h|-(1/h))} ds \\
&= 2e^{\pi/h} \int_0^1 |\phi_w(s)|^2 \left| \frac{2e^{-\pi|s/h|}}{|\phi_k(s/h)|^2} - 1 \right| e^{\pi(|s/h|-(1/h))} ds \\
&= 2h^{1+2\alpha} e^{\pi/h} \int_0^{1/h} \left| \frac{|\phi_w(1-hv)|}{(hv)^\alpha} \right|^2 \left| \frac{2e^{-\pi(1/h-v)}}{|\phi_k(1/h-v)|^2} - 1 \right| v^{2\alpha} e^{-\pi v} dv \\
&= 2h^{1+2\alpha} e^{\pi/h} o(1),
\end{aligned}$$

by the dominated convergence theorem. We have used the fact that both the functions $\phi_w(1-u)/u^\alpha$ and (see Lemma 5.1) $|(2 \exp(-\pi u)/|\phi_k(u)|^2) - 1|$ are bounded and that the second function is of order $O(1/u)$ as u tends to infinity. This shows that the term (5.4) is negligible with respect to (5.3). $\square$

Corollary 5.3. *The L^2 -norm of the function $\mathbf{v}_h$, defined in (3.5) is of order $O(h^{p(\frac{1}{2}+\alpha)} e^{p\pi/2h})$.*

Proof. This follows from the product form of $\mathbf{v}_h$ given by $\mathbf{v}_h(\mathbf{s}) = \prod_{j=1}^p v(s_j)$. $\square$

Proof of Theorem 4.1. The expansion (4.6) follows from Theorem 1 in Masry (1991). To prove the variance bound (4.7) we argue as in the proof of Theorem 2 in the same paper. First we give a bound on the variance in terms of the L_2 -norm of the function $\mathbf{v}_h$ and then we exploit the asymptotic expansion of the characteristic function ϕ_k as given in Lemma 5.1 to get a sharper bound on the L_2 -norm of $\mathbf{v}_h$ than Masry in his Proposition 3 by taking the behaviour of ϕ_w at the boundary of its support into account. Some details follow.

Argueing as in Masry (1991) we can show that

$$\text{Var } f_{nh}(\mathbf{x}) = O\left(\frac{\|\mathbf{v}_h\|_2^2}{nh^{2p}} + M_{nh}\right),$$

with (up to a multiplicative constant)

$$M_{nh} = \frac{1}{nh^{2p}} \sum_{j=p}^n \text{Cov}(W_j, W_0), \quad (5.5)$$

where $W_j = \mathbf{v}_h(\frac{\mathbf{x} - \log \mathbf{X}_j}{h})$.

Applying a lemma by Deo (1976), we can bound for strong mixing process X with mixing coefficients α_j the term M_{nh} by a constant (not depending on n and h) times

$$\frac{1}{nh^{2p}} \sum_{j=p}^n \alpha_{j-p+1}^\beta \{E |W_j|^{2/(1-\beta)} E |W_0|^{2/(1-\beta)}\}^{(1-\beta)/2},$$

which, by stationarity, becomes

$$\frac{1}{nh^{2p}} \sum_{j=p}^n \alpha_{j-p+1}^\beta (E |W_0|^{2/(1-\beta)})^{1-\beta}.$$

Observe now that, by boundedness of the density of $\log X_j^2$, the term $E |W_0|^{2/(1-\beta)}$ can be bounded by a constant times $h^p \|\mathbf{v}_h\|_{2/(1-\beta)}^{2/(1-\beta)}$ and that we can therefore write

$$\text{Var } f_{nh}(\mathbf{x}) = O\left(\frac{\|\mathbf{v}_h\|_2^2}{nh^p} + \sum_{j=p}^n \alpha_j^\beta \frac{\|\mathbf{v}_h\|_{2/(1-\beta)}^2}{nh^{p(1+\beta)}}\right).$$

The proof will be finished by application of Corollary 5.3, which gives the L^2 -norm of $\mathbf{v}_h$, and an estimate of the $L^{2/(1-\beta)}$ -norm of $\mathbf{v}_h$. For the latter one we have the inequalities $\|\mathbf{v}_h\|_{2/(1-\beta)} \leq \|\mathbf{v}_h\|_\infty^\beta \|\mathbf{v}_h\|_2^{1-\beta}$ and $\|\mathbf{v}_h\|_\infty \leq C \|\mathbf{v}_h\|_2$ for some constant C by the fact that ϕ_w has compact support. As a result we get $\|\mathbf{v}_h\|_{2/(1-\beta)} \leq C \|\mathbf{v}_h\|_2$ and that M_{nh} is less than a constant times $\|\mathbf{v}_h\|_2/nh^{p(1+\beta)}$. The bound on $\text{Var } f_{nh}(\mathbf{x})$ of theorem 4.1 now follows. $\square$

Proof of Theorem 4.2. Let $\mathcal{F}^\sigma$ be the σ -algebra generated by the process σ . We use the decomposition

$$\text{Var } f_{nh}(\mathbf{x}) = E \text{Var } (f_{nh}(\mathbf{x}) | \mathcal{F}^\sigma) + \text{Var } \tilde{f}_{nh}(\mathbf{x}), \quad (5.6)$$

with $\tilde{f}_{nh}(\mathbf{x})$ as in Remark 4.6. We now consider the first term in (5.6). Let $\mathbf{z}_j = (\log Z_j^2, \dots, \log Z_{j-p+1}^2)$ and $\mathbf{q}_j = (\log \sigma_j^2, \dots, \log \sigma_{j-p+1}^2)$. Since the Z_i are independent given $\mathcal{F}^\sigma$ we can bound the conditional variance by

$$\frac{1}{n^2 h^{2p}} \sum_{j=p}^n E \left[\left(\mathbf{v}_h \left(\frac{\mathbf{x} - \mathbf{q}_j - \mathbf{z}_j}{h} \right) \right)^2 | \mathcal{F}^\sigma \right]$$

which is by conditional independence and stationarity equal to

$$\frac{1}{nh^{2p}} \int \left(\mathbf{v}_h \left(\frac{\mathbf{x} - \mathbf{q}_0 - \mathbf{z}}{h} \right) \right)^2 k(\mathbf{z}) d\mathbf{z} \leq \frac{C}{nh^p} \|\mathbf{v}_h\|_2^2,$$

with C the maximum of $\mathbf{k}$, the density of $\mathbf{z}_0$. Therefore the first term in (5.6) is of order $\|\mathbf{v}_h\|_2^2/nh^p$, so of order $O(h^{p(1+2\alpha)} e^{p\pi/h}/nh^p)$.

The second term of (5.6) is treated next. We have with $U_j = \mathbf{w}(\frac{\mathbf{x}-\mathbf{q}_j}{h})$

$$\text{Var } \tilde{f}_{nh}(\mathbf{x}) = \frac{1}{n^2 h^{2p}} \sum_j \text{Var } U_j + \frac{2}{n^2 h^{2p}} \sum_{i < j} \text{Cov } (U_i, U_j).$$

The first term reduces by stationarity to $\frac{1}{nh^{2p}} \text{Var } U_1$ which can be bounded by a constant times $\|\mathbf{w}\|_2^2/nh^p$, since $(\log \sigma_1^2, \dots, \log \sigma_p^2)$ has by assumption a bounded density. For the second term we proceed as in the proof of Theorem 4.1. Using stationarity we write it as

$$\frac{2}{n^2 h^{2p}} \sum_{k=1}^n (n-k) \text{Cov } (U_k, U_0).$$

We split the summation into two parts. In the first part we consider

$$\sum_{k=1}^{p-1} (n-k) \text{Cov } (U_k, U_0).$$

whose absolute value can be bounded in view of the Cauchy-Schwarz inequality and stationarity by $(p-1)nE U_0^2$, which is bounded by $(p-1)nh^p \|\mathbf{w}\|_2^2$.

The absolute value of the second part

$$\sum_{k=p}^n (n-k) \text{Cov } (U_k, U_0)$$

can be bounded by invoking once more Deo's result by

$$n \sum_{k=p}^n \alpha_{k-p+1}^\beta (E |U_0|^{2/(1-\beta)})^{1-\beta},$$

which is less than

$$nh^{p(1-\beta)} \|\mathbf{w}\|_{2/(1-\beta)}^2 \sum_k \alpha_{k-p+1}^\beta.$$

Hence we have that $\text{Var } \tilde{f}_{nh}(\mathbf{x})$ is of order $1/nh^{p(1+\beta)}$.

Combining the obtained order estimates for the two terms of (5.6) and using the L^2 -norm of the function $\mathbf{v}_h$ gives the desired result. $\square$

References

- Abramowitz, M. and Stegun, I. (1964), *Handbook of Mathematical Functions, ninth edition*, Dover, New York.
- Bollerslev, T. (1986), Generalized autoregressive conditional heteroscedasticity, *J. Econometrics* **31**, 307–321.

- Bougerol, P. and N. Picard (1992a), Strict stationarity of generalized autoregressive processes, *Ann. Probab.* **20**, no. 4, 1714–1730.
- Bougerol, P. and N. Picard (1992b), Stationarity of GARCH processes and of some nonnegative time series, *J. Econometrics* **52**, no. 1-2, 115–127.
- Boussama, F. (1998), *Ergodicité, mélange et estimation dans les modèles GARCH*, PhD, thesis, Université Paris 7.
- Bradley, R.C. (1985), Basic properties of strong mixing conditions, in *Dependence in Probability and Statistics*, E. Eberlein and M.S. Taqqu Eds., Birkhäuser.
- Carasso, M. and Chen, X. (2002) Mixing and moment properties of various GARCH and stochastic volatility models, *Econometric Theory* **18**, 17–39.
- Deo, C.M. (1973), A note on empirical processes for strong mixing processes, *Ann. Probab.* **1**, 870–875.
- Doukhan, P. (1994), *Mixing, Properties and Examples*, Springer-Verlag.
- Fan, J. (1991), On the optimal rates of convergence for nonparametric deconvolution problems, *Ann. Statist.* **19**, 1257–1272.
- Hewitt, E. and Stromberg K. (1965), *Real and Abstract Analysis*, Springer Verlag, New York.
- Ibragimov, I.A., and Linnik (1971), *Independent and stationary sequences of random variables*, Wolters-Noordhoff.
- Masry, E. (1983), Probability density estimation from sampled data, *IEEE Trans. Inform. Theory* **29**, 696–709.
- Masry, E. (1991), Multivariate probability density deconvolution for stationary stochastic processes, *IEEE Trans. Inform. Theory* **37**, 1105–1115.
- Masry, E. (1993a), Asymptotic normality for deconvolution estimators of multivariate densities of stationary processes, *J. Mult. Anal.* **44**, 47–68.
- Masry, E. (1993b), Strong consistency and rates for deconvolution of multivariate densities of stationary processes, *Stoc. Proc. and Appl.* **475**, 53–74.
- Stefanski, L.A. (1990), Rates of convergence of some estimators in a class of deconvolution problems, *Statist. Probab. Lett.* **9**, 229–235.

- De Vries, C.G. (1991), On the relation between GARCH and stable processes, *J. Econometrics* **48**, 313–324.
- Wand, M.P.. (1998), Finite sample performance of deconvolving kernel density estimators, *Statist. Probab. Lett.* **37**, 131–139.
- Wand, M.P. and Jones, M.C. (1995), Kernel Smoothing, Chapman and Hall, London.